

Abstract Submitted  
for the Thirteenth International Conference on Laser  
Interactions and Related Plasma Phenomena (LIRPP)  
April 13-18, 1997

Semiclassical Atomic Model.\* R. M. MORE, J. LEVATIN,  
Lawrence Livermore National Laboratory, T. KATO(\*\*) —  
Simulation of laser-matter interactions requires a broad range  
of atomic data for nonequilibrium (NTLE) atomic processes,  
equations of state, photon absorption opacity and charged-  
particle stopping powers. There is a continuing need for a  
simple and flexible source of atomic data which could improve  
the results given by the screened hydrogenic model without  
requiring too much computer time. We have developed  
techniques for working with WKB wave functions including  
relativistic corrections and calculation of matrix elements for  
radiative transitions. Here we assemble these results in a  
semiclassical j-j coupled relativistic self-consistent field model.  
This model obeys various consistency tests and is embodied in  
a simple and robust computer code.

\*Supported in part under the auspices of the U.S. Department  
of Energy by Lawrence Livermore National Laboratory under  
Contract No. W-7405-ENG-48.

(\*\*)NIFS, Nagoya, Japan; supported by U.S.-Japan  
Cooperation Program (Japan Society for the Promotion of  
Science).

Submitted by

Richard M. More  
Lawrence Livermore National Lab  
P. O. Box 808, L-041  
Livermore, CA 94550